



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Fracture healing and the underexposed role of extracellular vesicle-based crosstalk

Qiao, Zhi ; Greven, Johannes ; Horst, Klemens ; Pfeifer, Roman ; Kobbe, Philipp ; Pape,
Hans-Christoph ; Hildebrand, Frank

Abstract: The process of fracture healing is complex and requires an interaction of multiple organ systems. Cell-cell communication is known to be very important during this process. Extracellular vesicle (EVs) are small membranous vesicles generated from a variety of cells. Proteins, RNAs, small molecules and mitochondria DNA were found to be transported among cells through EVs. EV-based crosstalk represents a substantial cell-cell communication pattern, that can both interact with cells through molecular surfaces, and transfer molecules to cells. These interactions can assist in the synchronization of cellular functions among cells of the same kind, and coordinate the functions of different types of cells. After activation, platelets, neutrophils, macrophages, osteoblasts, osteoclasts, and mesenchymal stem cell (MSC) all secrete EVs, promoting the fracture healing process. Moreover, some studies have found evidence that EVs may be used for diagnosis and treatment of delayed fracture healing, and may be significantly involved in the pathophysiology of fracture healing disturbances. In this review, we summarize recent findings on: 1) EVs released by fracture healing-related cells, and 2) EV-mediated communications during fracture healing. We also highlight the potential applications of EVs in fracture healing. Lastly, the prospect of EVs for research and clinical use is discussed.

DOI: <https://doi.org/10.1097/SHK.0000000000001002>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-139864>

Journal Article

Published Version

Originally published at:

Qiao, Zhi; Greven, Johannes; Horst, Klemens; Pfeifer, Roman; Kobbe, Philipp; Pape, Hans-Christoph; Hildebrand, Frank (2018). Fracture healing and the underexposed role of extracellular vesicle-based crosstalk. *Shock*, 49(5):486-496.

DOI: <https://doi.org/10.1097/SHK.0000000000001002>

Fracture healing and the underexposed role of extracellular vesicle-based crosstalk

Zhi Qiao¹, Johannes Greven¹, Klemens Horst¹, Roman Pfeifer², Philipp Kobbe¹, Hans-Christoph Pape²,
Frank Hildebrand¹

Affiliations:

1. Department of Trauma and Reconstructive Surgery RWTH Aachen University Hospital
2. Department of Orthopaedic Trauma and Harald-Tscherne Laboratory, University Hospital Zurich, University of Zurich; Ramistr. 100, 8091 Zuerich, Switzerland

Corresponding author:

Zhi Qiao

Department of Trauma and Reconstructive Surgery RWTH Aachen University Hospital

Pauwelsstreet 30

52074 Aachen

Germany

Tel: 0049 241 89117

Qiaozhi_1988@hotmail.com

Conflict of interest:

Zhi Qiao is supported by the *China Scholarship council* (No. 201508080049)

Running head:

extracellular vesicle-based crosstalk in fracture healing

Acknowledgements:

This article was proofread by *Proof-Reading-Service.com*

Abstract

The process of fracture healing is complex and requires an interaction of multiple organ systems. Cell-cell communication is known to be very important during this process. Extracellular vesicle (EVs) are small membranous vesicles generated from a variety of cells. Proteins, RNAs, small molecules and mitochondria DNA were found to be transported among cells through EVs. EV-based crosstalk represents a substantial cell-cell communication pattern, that can both interact with cells through molecular surfaces, and transfer molecules to cells. These interactions can assist in the synchronization of cellular functions among cells of the same kind, and coordinate the functions of different types of cells. After activation, platelets, neutrophils, macrophages, osteoblasts, osteoclasts, and mesenchymal stem cell (MSC) all secrete EVs, promoting the fracture healing process. Moreover, some studies have found evidence that EVs may be used for diagnosis and treatment of delayed fracture healing, and may be significantly involved in the pathophysiology of fracture healing disturbances. In this review, we summarize recent findings on: 1) EVs released by fracture healing-related cells, and 2) EV-mediated communications during fracture healing. We also highlight the potential applications of EVs in fracture healing. Lastly, the prospect of EVs for research and clinical use is discussed.

Keywords

Extracellular vesicle; Osteoblast; Osteoclast; Platelet; Endothelial cell; Intercellular communication; Crosstalk; MiRNA

Fracture healing is a complex process, involving the interaction of immunity, bone, and vessel systems. During bone healing, cells from different systems have to communicate with each other. There are three cell-cell signal transduction modes that are commonly known and accepted(1):

- 1) secretion molecules,
- 2) cell-cell contact, and
- 3) tunneling nanotubes.

Recent studies have introduced the potential for communication via extracellular vesicles (EVs)(2, 3). It is known that EVs from osteoblasts, chondroblasts, and osteoclasts are important for tissue calcification and bone remodeling. Furthermore, studies have shown a crucial role of EV-based crosstalk in the physiological and pathophysiological processes of coagulation, inflammation, and osteogenesis(3, 4). These processes are involved in fracture healing, and thus, EV-based crosstalk communication occurs throughout the whole process of fracture healing. As different types of EVs can be released under different conditions(5, 6), this review focuses on studies dealing with the role of EVs in the process of fracture healing.

1. What are EVs

EVs are membranous vesicles generated from a variety of cells, with diameters ranging from 30-5000 nm. Three kinds of extracellular EVs have been observed: exosomes, shedding microvesicles, and apoptotic blebs(7). Exosomes are cup-shaped membranous EVs, with a diameter of 30-100 nm. They are generated by inward budding of endosomal membranes into large multivesicular bodies (MVBs), and secreted by exocytosis(8), with surface markers like Alix, HSC70, CD63, CD81, and CD9 (7). Shedding microvesicles are generated by blebbing and shedding of the plasma membrane from almost all cell types, with a diameter of 100-1000 nm, and have cell markers such as selectins and integrins(9). In some publications, shedding microvesicles were also called microparticles(10), while in some others, microparticles referred as those isolated from platelets or endothelial cells(9). Apoptotic blebs are membrane vesicles generated by apoptotic or dying cells(7), with a diameter of 50-5000 nm. Inside

of these EVs, proteins, RNAs, small molecules and mitochondria DNA can be found. In this article, we focus on exosomes and shedding microvesicles.

2. EV-based crosstalk

EVs are competent information mediators. The membranous structure of EVs enables them to transport molecules within their inner cytosol, and membrane-dependent molecules at the surface. The specialties of EV-based molecular transport are:

- 1) Transport of RNAs between cells,
- 2) Transport of components of specific pathways
- 3) Transport of surface molecules and membranous molecules.

The transportation of these molecules enables them to coordinate cellular reactions; this process can control gene expression by regulating mRNA turnover, ultimately contributing to the epigenetic and proteomic properties of the target cells (11), as well as changing the maturation and differentiation of target cells (12). At the same time, cells can activate surface molecular pathways of remote cells.

Organelles have also been found to be transferred between different cells by EVs (13). Studies have found mitochondrial DNAs in EVs from platelets, astrocytes, glioblastoma, and bone marrow-derived stromal cells (14, 15). According to the results from recent studies, the release of mitochondria is able to elevate inflammatory reactions (14, 16).

In addition to other mechanisms, the interaction of EVs with cells includes 1) surface molecules, 2) EV-cell fusion, and 3) cargo releasing.

EVs carry surface molecules and transmembrane proteins on their surfaces and, therefore, include characteristics of the cells they were released from. These surface proteins help to interact with specific target cells, increase the specificity of cargo delivery and trigger signaling cascades via receptor interactions(7).

Through EV-cell fusion, proteins and RNAs can be transported into recipient cells. The EV-cell interactions and fusions are mediated by receptor binding (17). Using membrane-bound structures, EVs can receive signals and fuse with their target cells(3). Due to the presence of surface receptors, EVs only interact with cells they recognize(18).

EVs also have the potential to reach the extracellular spaces and blood, releasing their content and membrane components(18, 19). EVs contain proteinase, cytokines, growth factors, and organelles(1, 20) that function in the extracellular matrix.

3. EVs from fracture healing-related cells

3.1 Osteoblastic EVs (OBEVs)

Osteoblasts play a central role in the fracture healing process. It has already been reported that osteoblasts directly communicate with osteoclasts (4, 21), mesenchymal stem cell (MSC)(22, 23), and others (24), via EVs. Proteomic analysis of OBEVs/exosomes revealed (25) proteins that modulate osteogenesis, which are involved in pathways such as the eukaryotic initiation factor 2 (EIF2) signaling pathway, integrin signaling pathway, Rho-guanosine triphosphate (Rho-GTP) signaling pathway, and mammalian target of rapamycin (mTOR) signaling pathway(25).

OBEVs were observed to induce differentiation of multi-potential stem cells into osteoblasts (24, 26, 27). OBEVs/exosomes from mineralizing osteoblasts have the ability to promote the differentiation of bone marrow stromal cells into osteoblasts, a process which is mediated by the Wnt signaling pathway(27). Several studies have also found evidence to show that osteoblasts communicate with osteoclasts through OBEVs. Receptor activator of nuclear factor- κ B ligand (RANKL) proteins, which

are able to induce generation of osteoclasts, were found in EVs/microvesicles originated from osteoblasts(4). In rats, hypoxia stimulates the release of EVs/microvesicles from osteoblasts, containing vesicular adenosine triphosphate (ATP), which was proved to influence functions of osteoclasts(28).

3.2 Mesenchymal stem cell(MSC)-derived EVs (MSCEV)

MSCs play an active role in the processes of immune modulation and tissue regeneration. Cytokines, micro-RNAs (miRNAs), surface molecules, and signaling pathway components from MSCEV have been shown to be involved in cell-cell communications.

Ischemia-activated MSCs can release EVs with high concentrations of platelet-derived growth factor (PDGF), epidermal growth factor (EGF), fibroblast growth factor (FGF), and other angiogenesis-related molecules(29). The release of these factors into the extracellular matrix promotes angiogenesis and osteogenesis.

Important fracture healing-related pathways have been found in EVs/microvesicles from activated MSCs, including in the Wnt signaling pathway, mitogen-activated protein kinase (MAPK) signaling pathway, transforming growth factor (TGF) signaling pathway, peroxisome proliferator-activated receptors (PPAR) signaling pathway, and the bone morphogenetic protein (BMP) signaling pathway(30). A variety of surface molecules from MSCEVs/microvesicles were found to be involved in fracture healing processes, including PDGF receptors, EGF receptors, ezrin (EZR), integrins, and IQ motif containing GTPase activating protein 1 (IQGAP-1) (30). These pathways and molecules are involved in the recruitment and differentiation of MSCs(31), the regulation of endothelial cell proliferation, angiogenesis(32), as well as epithelial proliferation(33).

A variety of miRNAs have been detected in MSCEVs/microvesicles(34), some of which were also shown to have functions related to fracture healing. Nakamura et al. (35) reported that MSC-derived miRNA promotes myogenesis and angiogenesis. MiR-196a(22) was observed to promote osteogenesis, while miR-378(36), miR-223(37), and miR-100(38) were found to inhibit osteogenesis. MiR-378(39) and miR-223(40) were also found to be involved in angiogenic remodeling. It is also reported

that, at different time points during fracture healing, MSCs release EVs with different profiles of miRNAs, which are involved in osteoblastogenesis or leukocyte migration (23)

3.3 Endothelial EVs (EEVs)

The generation of endothelial EVs was found to be stimulated by inflammatory cytokines, bacterial lipopolysaccharides, reactive oxygen species, thrombin, C-reactive protein, and uremic toxins (41). The generation of EEVs after TNF- α and thrombin stimulation involves several signaling pathways including the caspase, rho-kinase, nuclear factor kappa B (NF- κ B) pathways, and MAPK (41, 42). Tumor Necrosis Factor α (TNF- α), interleukin -1 (IL-1), and integrin-1 may amplify the release of EEVs through the NF- κ B and MAPK pathways (43).

EEVs have been found to be important elements in the processes of coagulation, inflammation, angiogenesis, and osteogenesis. In the early phase of fracture healing, phospholipids and proteins at the surface of EEVs were found to control the coagulation equilibrium, among them are endothelial protein C receptor (EPCR), thrombomodulin (TM), tissue factor (TF), von Willebrand factor (vWf), tissue plasminogen activator (tPA), and Phosphatidylserines (PS) (43-46). Inflammation-activated EEVs bear inflammation modulating molecules such as platelet endothelial cell adhesion molecule-1 (PECAM-1/CD31), intercellular adhesion molecule 1 (ICAM-1), vascular cell adhesion molecule (VCAM-1), and endothelial-selectin (E-selectin) (42, 47). These molecules are known to have diverse roles in vascular biology, including angiogenesis, thrombosis, and regulation of multiple stages of leukocyte migration. At the same time, studies have reported that EEVs modulate proliferation, angiogenesis, and apoptosis of endothelial cells through RNA transportation. Horizontal transportation of miRNAs between endothelial cells by EEVs/microvesicles promotes angiogenesis following Akt activation and endothelial nitric oxide synthase (eNOS) expression (48), and promotes vascular endothelial repair (49). Furthermore, EEVs contain matrix metalloproteases, which are involved in the extracellular matrix degradation, as well as the release of growth factors that play crucial roles in bone remodeling and angiogenesis (50, 51). The role of EEVs in osteogenesis has been reported in several studies. Inflammation-induced EEVs express molecules, such as PECAM-1 and ICAM-1, which were found to promote osteoclastogenesis (52, 53). EEVs/microvesicles from TNF- α -stimulated endothelial cells were also found to contain a significant amount of BMP-2, and were able to enhance vascular smooth muscle cell osteogenesis and calcification (54).

3.4 OsteoclasticEVs (OCEVs)

Osteoclasts are an important element in fracture healing and bone remodeling. A study tested the profile of OCEV-miRNAs from TNF- α and RANKL-stimulated osteoclasts, showing a variety of miRNAs (miR-378, miR-223, mir-210, miR-21, miR-155, miR-146a, and miR-199a) that may be involved in angiogenesis, osteogenesis, and chondrogenesis(55). RANK was also found to be present on the surface of osteoclasticEVs, and is capable of regulating osteoclastogenesis(56). More studies are needed to investigate the communication functions of OCEVs.

3.5 Macrophage and monocyte-derivedEVs

Macrophages and monocytes are important for both initiation of inflammatory reactions and tissue repair. Granulocyte-macrophage colony stimulating factor (GM-CSF)(57), Lipopolysaccharides (LPS)(58), and Fas-Ligand(59) were found to be able to stimulate generation of EVs by macrophages.

EVs/microvesicles from cultured macrophages have been shown to transport functional RNA molecules into target cells, including monocytes, endothelial cells, epithelial cells, and fibroblasts(57). EVs/microvesicles from monocytes and macrophages were found to be rich of tissue factor (TF), which can bind platelets through P-selectin glycoprotein ligand-1 (PSGL-1) on the EVs, and P-selectin on the platelets; thereby enhancing coagulation(60). EVs/microvesicles from monocytes were observed to interact with endothelial cells, inducing pro-coagulatory and pro-inflammatory endothelial phenotypes(59). *In vitro* capillary tube formation and *in vivo* angiogenesis assays have shown that monocyte-derived EVs/microvesicles have strong pro-angiogenic activities(61). MSCs/exosomes incubated by monocyte-derived EVs showed increased runt-related transcription factor 2 (RUNX2) and BMP-2, indicating the osteogenic differentiation of MSCs(58).

3.6 Neutrophil-derivedEVs (NEVs)

The initiation of an inflammatory and coagulatory response has the potential to stimulate the generation of NEVs. TNF- α , the complement system, IL-8, platelet activating factor protein kinase C activator, and nitric oxide synthase inhibitor have been found to increase the production of NEVs(62). It has been reported that, under inflammatory conditions, the amount of NEVs will increase significantly, compared to normal conditions(63, 64).

Although there are no studies referring to the functions of NEVs in fracture healing, several studies have indicated a role of NEVs in fracture healing-related reactions. During coagulation, platelets were found to be activated by NEVs in an Akt phosphorylation-dependent manner(65). In addition, NEVs appear to be important inflammation modulators, as they may have both anti-inflammatory(10, 66) and pro-inflammatory functions(10, 67, 68), depending on the background of the microenvironment(10, 62). Studies on proteomes of NEVs/microvesicles have revealed over 400 proteins, covering mainly integrin signaling, EGF, and FGF signaling pathways(10). Although it has not been proven yet, the EGF and FGF signaling pathways found in NEVs/microvesicles may stimulate the differentiation of MSCs and angiogenesis(10).

3.7 Platelet-derived EVs (PEVs)

Platelets are activated under coagulatory and inflammatory conditions, with the subsequent release of large amounts of functional PEVs(69). A study demonstrated that 87% of the release of platelet-activating factor (PAF), a key factor for triggering inflammatory and thrombotic cascades, was associated with EVs(70, 71). There is also evidence that PEVs may transfer platelet-specific immunoreactive antigens to the surface of other cells, modulating cellular functions(72).

In the early phase of fracture healing, expansion of coagulation platelet-platelet crosstalk is one of the most important features of PEVs. PEVs express anionic phospholipids (AP) and TF on their surfaces; arachidonic acids (AA) from platelets were reported to have the potential to activate the clotting cascades in platelets(73, 74). PEVs have also been found to be able to bind to the subendothelial matrix *in vitro* and *in vivo*, acting as substrates for further platelet binding(75).

Posttraumatic modulation of inflammation by PEV-based crosstalk was shown in several publications(76). In a clinical study, high posttraumatic levels of PEVs were associated with the development of acute respiratory distress syndrome (ARDS) and multiple organ failure(MOF)(73). Through PEV AA and bioactive prostanoids, PEVs were found to promote inflammatory reactions by activating MAPK pathways(77). Through the MAPK pathway, PEVs can induce the production of

important inflammation mediators like Cyclooxygenase-2 (COX-2) and prostacyclin (PGI₂), in human endothelial cells(77). PEVs were shown to activate monocytes and macrophages, and reprogram their functions towards a phagocytic phenotype, by regulating their RNA expression(78) and activating the NF- κ B pathway(79). Neutrophils(80) and leukocytes(16, 81) are also the targets of PEVs. One study found that the binding of PEVs to neutrophils can induce a significant increase in phagocytic activity in a concentration-dependent manner(80). Mitochondria found in PEVs can perform as inflammation mediators, activating leukocytes and neutrophils, causing inflammatory responses(16).

Studies have shown that PEVs can modulate the proliferation, survival, chemotaxis, and function of endothelial cells through the VEGF pathway(82) and by argonaute 2 (Ago2)-microRNA complex-delivering(83). It has also been demonstrated that PEVs are able to stimulate mitogenic activity of bone cells, contributing to bone regeneration(84). In addition, a study showed that surface molecules on PEVs can activate hematopoietic cells by chemotaxis, increasing adhesion, proliferation, and survival, and activating signaling cascades including MAPK p42/44, phosphatidylinositol-3 kinase (PI-3K), AKT, and Signal transducer and activator of transcription (STAT) proteins(85).

3.8 Other cellular EVs involved in fracture healing processes

Other cell types have also been observed to release EVs under inflammatory conditions. Even though there is limited evidence in the literature, the existing data may provide the first evidence for the role of EV-based crosstalk in fracture healing. Erythrocytes activated by lysophosphatidic acid can secrete pro-coagulatory EVs, which contribute to coagulation processes(86). Vascular smooth muscle cells incubated by LDL are found to produce TF-bearing EVs, promoting coagulation cascades(87).

EVs from chondrocytes have been detected in the growth plate. EVs are released in a polarized fashion from the lateral edges of growth plate chondrocytes. They have been found to be involved in the mineralization of bone matrix(88). Later, BMP, VEGF, bone sialoprotein (BSP), osteonectin, and osteocalcin were also found in these EVs, highlighting the communicative role of chondroblastic EVs(88-91). VEGF, if released into the matrix, regulates angiogenesis and osteogenesis, and stimulates endothelial cells, chondroblasts, and osteoblasts(92). BSP in EVs was shown to promote differentiation of osteoblasts and repair of bone defects(91), as well as to promote differentiation of

chondrocytes and osteoblasts(in growth periods) (90). Further, osteopontin, osteonectin, and osteocalcin were found to be able to inhibit mineralization *in vitro*(88).

4. EV-based crosstalk network during fracture healing

Fracture healing involves very complex systemic reactions. It involves sophisticated cooperation between a variety of cells, cytokines, and inflammatory factors (figure 3). In this chapter, we will briefly overview the network of EV-based crosstalk during fracture healing, highlighting the networks of cytokines and inflammatory factors.

Coagulation and inflammatory reactions are the first steps for fracture healing. In this phase, the ruptured blood vessels, tissue necrosis, and other ongoing damage, stimulate the inflammatory cascades. The activation of platelets, neutrophils, and macrophages by hypoxia, coagulation, and inflammation are observed to produce, not only a variety of cytokines and inflammatory factors, but also large amounts of active EVs(1, 93). The appearance of EVs may indicate that they have important functions.

A short time after blood vessel and tissue injury, pro-coagulatory EVs can be released from platelets and neutrophils. The positive feedback model of PEVs appears to further stimulate PEV productions (74), as well as activate and attract neutrophils(16, 80) and macrophages(78, 79). EEVs from activated endothelial cells also promote coagulation via EPCR, TM, TF, vWf, tPA, PS, and plasminogen(43, 94, 95).

Inflammatory background is important for fracture healing. EVs derived from a variety of cells participate in the inflammatory regulation of the fracture site. PEVs are able to stimulate the differentiation of monocytes and macrophages, inducing the production of pro-inflammatory factors(78, 79). The activation of neutrophils results in secretion of NEVs, that also regulate inflammatory reactions. In addition, NEVs can increase the inflammatory response by the activation and

aggregation of macrophages(67), inducing production of IL-6, IL-8, MAC-1, and oxygen active species(10, 96). However, NEVs can also inhibit the aggregation of neutrophils (97), induce the production of anti-inflammatory factors from macrophages such as IL-10 and TGF- β 1(66, 98) and inhibit pro-inflammatory actions of endothelial cells(10). Although EVs from MSCs have been shown to have the potential to modulate the inflammatory response, further studies are needed to clarify their role in the process of fracture healing.

A variety of EVs take part in angiogenesis modulation during fracture healing. EVs from platelets, neutrophils, macrophages, MSCs, and endothelial cells have been shown to have the ability to stimulate differentiation of stem cells or endothelial cells. EVs from MSCs(29, 31) and chondrocytes(92) have been shown to release multiple growth factors, providing a microenvironment for angiogenesis and tissue regeneration. Surface molecules from MSCEVs(32, 33) and EEVs(99) have also been shown to modulate angiogenesis. Pathway molecules and miRNAs from PEVs(83), MSCEVs(35), EEVs(48, 49, 100), and osteoclast-(101) and monocyte-derived EVs(61) also participate in the modulation of angiogenesis.

During tissue regeneration and bone remodeling, multiple EVs take part in the process of activation of fibroblasts, chondroblasts, and osteoblasts. EVs from MSCs have a significant role in wound healing. They were found to enhance wound healing via promotion of collagen synthesis and angiogenesis(102). One study also found that MSC-derived exosomes caused an increase in proliferation and migration of fibroblasts(103). In the microenvironment of osteogenesis, a mixture of EVs has been found to modulate the osteogenesis reactions. Matrix vesicles found in the growth plate serve as carriers of morphogenetic information to nearby chondrocytes and osteoblasts. These matrix vesicles were observed to be released from the lateral edges of growth plate chondrocytes, the osteoid-facing surfaces of osteoblasts, and the apical surfaces of odontoblasts(88). Cultured chondrocytes treated with these matrix vesicles showed a two- to three-fold increase in alkaline phosphatase activity (88). OBEVs contain miRNAs and pathway molecules, which have the potential to stimulate osteogenesis of MSCs(26, 27), and communicate with osteoclasts for bone remodeling(4, 28). Molecules from EEVs, such as BMP-2(54) and PECAM-1(104) have been shown to involve in the modulation of angiogenesis, osteogenesis, and bone remodeling.

In addition to their communicative functions, EVs from osteoblasts, chondroblasts, and osteoclasts are also involved in osteogenesis and bone remodeling. They are the practical workers for bone matrix generation and calcification. Aspects of matrix generation, calcification, and bone remodeling are reviewed elsewhere(105).

A summary of EV-based miRNAs related to osteogenesis, obtained from ExoCarta (12/2016)(106), can be found in Figure 1, Figure2, and Table 1.

5. The potential application value of EVs during fracture healing

5.1 Molecular diagnosis

Multiple components of EVs, especially miRNAs, are suggested as biomarkers for molecular diagnosis. Several studies have discussed the diagnostic value of EVs in neurological disorders(107), cardiovascular diseases(108), osteoporosis(48)and sepsis(109).

With regards to the skeleton, the diagnostic potentials have also been investigated. One study tested the proportion of EVs in rheumatoid arthritis (RA) patients. The authors found an increase of plasmatic CD3(+), CD14(+), CD19(+), CD41(+), CD62E(+) MPs, and urinary CD14(+), CD3(+), and CD19(+) MPs in RA patients with high disease activity(110). More interestingly, compared to a healthy population, the level of miR-92a was found to decrease 24 hours after a bone fracture, while it was decreased 14 days after a lumbar compression fracture(111). These studies showed the potential of EVs as a diagnostic tool. However, more research is needed to bring this marker into clinical practice as a routine parameter.

5.2 EV-based therapies

Comparedto proteins and RNAs, EV structures may be more tolerant to the microenvironment *in vivo*(112). Thus, the use of EEV-mediated delivery is expected to be more efficient than using soluble

proteins or RNA molecules. One study also indicated that EV structures may be at lower risk of immunogenicity and toxicity(112).

The therapeutic potential of EVs has been studied in the treatment of several diseases, such as acute lung injury(113), myocardial ischemia/reperfusion injury (114), kidney injury (115), peri-implant bone defect(116) and others; the results appear to be promising. Moreover, some researchers have developed manufactured EVs to improve treatment of fracture healing. In this context, researchers aimed to remodel cell-derived EVs, thereby making them more effective for treatment. These small molecules, miRNAs, and exogenous proteins have been successfully delivered by EVs(117, 118). In addition, specifically generated gelatin EVs have been used as vehicles to transport molecules. With this technique, bone defects have been treated successfully by transporting BMPs, VEGFs, TGF- β , and other molecules(119, 120), to target cells. Poly(lactic-co-glycolic acid) (PLGA) nanoparticles were also shown to be a potential vehicle in guiding bone regeneration(121).

6. Prospect for research and clinical use

To date, the mechanism of EVs involvement in bone matrix generation and mineralization has been thoroughly investigated. However, there is still little information about the role of information transfer during the process of fracture healing. Therefore, further studies should focus on this area of research. To date, knowledge gained from studies has only shown the important role of EV-based communication in fracture healing, while the more detailed mechanisms remain unknown.

Many potential advantages of EVs make them worthy for future studies. EVs can be easily collected from blood or body fluids, with comparable characteristics of their original solid cells; this is advantageous for the diagnostic application of EVs. MiRNAs and surface molecules in EVs are more stable. However, little is known about the specificity of EVs in fracture healing-related states. Further studies are needed to look into the components and surface markers of EVs which may be able to predict the fracture healing states, and the potential threats.

The application of EVs in tissue engineering offers many possibilities for treatment options. Researchers have tried to use the advantages of EVs in improving fracture healing and treating bone defects, with optimistic results. However, some restrictions limit their large scale clinical use. For example, the production of EVs should be generated with more specific components, and needs to be produced in large scale. Thus, specific tissue-engineering cells may be a good option for the next research step. Tissue-engineering cells or bacteria can produce highly concentrated and purified EVs. At the same time, highly specified surface molecules can accurately aim at targeted cells. In addition, the clearance of EVs *in vivo* also needs to be considered; more stable EVs may need to be developed for effective treatments. Furthermore, man-made EVs represent a very promising field for research and treatment.

7. Conclusion

EVs play an important role during fracture healing, with their information transportation and bone matrix generation. While the mechanism of EV involvement in bone matrix generation is clear, the mechanism of EVs role in information transportation remains underexposed and is a field of research which requires further study. It is likely that, in the near future, EVs will be useful in clinical diagnosis and treatment.

Reference

1. Ratajczak J, Wysoczynski M, Hayek F, Janowska-Wieczorek A and Ratajczak MZ: Membrane-derived microvesicles: important and underappreciated mediators of cell-to-cell communication. *Leukemia* 20(9):1487-95, 2006.
2. Hugel B, Martinez MC, Kunzelmann C and Freyssinet JM: Membrane microparticles: two sides of the coin. *Physiology (Bethesda)* 20:22-7, 2005.
3. Cocucci E, Racchetti G and Meldolesi J: Shedding microvesicles: artefacts no more. *Trends Cell Biol* 19(2):43-51, 2009.
4. Deng L, Wang Y, Peng Y, Wu Y, Ding Y, Jiang Y, Shen Z and Fu Q: Osteoblast-derived microvesicles: a novel mechanism for communication between osteoblasts and osteoclasts. *Bone*, 2015.
5. Peterson DB, Sander T, Kaul S, Wakim BT, Halligan B, Twigger S, Pritchard KA, Jr., Oldham KT and Ou JS: Comparative proteomic analysis of PAI-1 and TNF-alpha-derived endothelial microparticles. *Proteomics* 8(12):2430-46, 2008.
6. Milioli M, Ibanez-Vea M, Sidoli S, Palmisano G, Careri M and Larsen MR: Quantitative proteomics analysis of platelet-derived microparticles reveals distinct protein signatures when stimulated by different physiological agonists. *J Proteomics* 121:56-66, 2015.
7. Mathivanan S, Ji H and Simpson RJ: Exosomes: extracellular organelles important in intercellular communication. *J Proteomics* 73(10):1907-20, 2010.
8. Simpson RJ, Lim JW, Moritz RL and Mathivanan S: Exosomes: proteomic insights and diagnostic potential. *Expert Rev Proteomics* 6(3):267-83, 2009.
9. Lawson C, Vicencio JM, Yellon DM and Davidson SM: Microvesicles and exosomes: new players in metabolic and cardiovascular disease. *J Endocrinol* 228(2):R57-71, 2016.
10. Dalli J, Montero-Melendez T, Norling LV, Yin X, Hinds C, Haskard D, Mayr M and Perretti M: Heterogeneity in neutrophil microparticles reveals distinct proteome and functional properties. *Mol Cell Proteomics* 12(8):2205-19, 2013.
11. Hunter MP, Ismail N, Zhang X, Aguda BD, Lee EJ, Yu L, Xiao T, Schafer J, Lee ML, Schmittgen TD, et al.: Detection of microRNA expression in human peripheral blood microvesicles. *PLoS One* 3(11):e3694, 2008.
12. Eken C, Gasser O, Zenhausem G, Oehri I, Hess C and Schifferli JA: Polymorphonuclear neutrophil-derived ectosomes interfere with the maturation of monocyte-derived dendritic cells. *J Immunol* 180(2):817-24, 2008.
13. Spees JL, Olson SD, Whitney MJ and Prockop DJ: Mitochondrial transfer between cells can rescue aerobic respiration. *Proc Natl Acad Sci U S A* 103(5):1283-8, 2006.
14. Islam MN, Das SR, Emin MT, Wei M, Sun L, Westphalen K, Rowlands DJ, Quadri SK, Bhattacharya S and Bhattacharya J: Mitochondrial transfer from bone-marrow-derived stromal cells to pulmonary alveoli protects against acute lung injury. *Nat Med* 18(5):759-65, 2012.
15. Guescini M, Genedani S, Stocchi V and Agnati LF: Astrocytes and Glioblastoma cells release exosomes carrying mtDNA. *J Neural Transm (Vienna)* 117(1):1-4, 2010.
16. Boudreau LH, Duchez AC, Cloutier N, Soulet D, Martin N, Bollinger J, Pare A, Rousseau M, Naika GS, Levesque T, et al.: Platelets release mitochondria serving as substrate for bactericidal group IIA-secreted phospholipase A2 to promote inflammation. *Blood* 124(14):2173-83, 2014.

17. Losche W, Scholz T, Temmler U, Oberle V and Claus RA: Platelet-derived microvesicles transfer tissue factor to monocytes but not to neutrophils. *Platelets* 15(2):109-15, 2004.
18. Gasser O and Schifferli JA: Activated polymorphonuclear neutrophils disseminate anti-inflammatory microparticles by ectocytosis. *Blood* 104(8):2543-8, 2004.
19. Mochizuki S and Okada Y: ADAMs in cancer cell proliferation and progression. *Cancer Sci* 98(5):621-8, 2007.
20. Moldovan L, Batte K, Wang Y, Wisler J and Piper M: Analyzing the circulating microRNAs in exosomes/extracellular vesicles from serum or plasma by qRT-PCR. *Methods Mol Biol* 1024:129-45, 2013.
21. Zhao H: Membrane trafficking in osteoblasts and osteoclasts: new avenues for understanding and treating skeletal diseases. *Traffic* 13(10):1307-14, 2012.
22. Qin Y, Wang L, Gao Z, Chen G and Zhang C: Bone marrow stromal/stem cell-derived extracellular vesicles regulate osteoblast activity and differentiation in vitro and promote bone regeneration in vivo. *Sci Rep* 6:21961, 2016.
23. Xu JF, Yang GH, Pan XH, Zhang SJ, Zhao C, Qiu BS, Gu HF, Hong JF, Cao L, Chen Y, et al.: Altered microRNA expression profile in exosomes during osteogenic differentiation of human bone marrow-derived mesenchymal stem cells. *PLoS One* 9(12):e114627, 2014.
24. Nair R, Santos L, Awasthi S, von Erlach T, Chow LW, Bertazzo S and Stevens MM: Extracellular vesicles derived from preosteoblasts influence embryonic stem cell differentiation. *Stem Cells Dev* 23(14):1625-35, 2014.
25. Ge M, Ke R, Cai T, Yang J and Mu X: Identification and proteomic analysis of osteoblast-derived exosomes. *Biochem Biophys Res Commun* 467(1):27-32, 2015.
26. Narayanan R, Huang CC and Ravindran S: Hijacking the Cellular Mail: Exosome Mediated Differentiation of Mesenchymal Stem Cells. *Stem Cells Int* 2016:3808674, 2016.
27. Cui Y, Luan J, Li H, Zhou X and Han J: Exosomes derived from mineralizing osteoblasts promote ST2 cell osteogenic differentiation by alteration of microRNA expression. *FEBS Lett* 590(1):185-92, 2016.
28. Orriss IR, Knight GE, Utting JC, Taylor SE, Burnstock G and Arnett TR: Hypoxia stimulates vesicular ATP release from rat osteoblasts. *J Cell Physiol* 220(1):155-62, 2009.
29. Anderson JD, Johansson HJ, Graham CS, Vesterlund M, Pham MT, Bramlett CS, Montgomery EN, Mellema MS, Bardini RL, Contreras Z, et al.: Comprehensive Proteomic Analysis of Mesenchymal Stem Cell Exosomes Reveals Modulation of Angiogenesis via NFkB Signaling. *Stem Cells*, 2016.
30. Kim HS, Choi DY, Yun SJ, Choi SM, Kang JW, Jung JW, Hwang D, Kim KP and Kim DW: Proteomic analysis of microvesicles derived from human mesenchymal stem cells. *J Proteome Res* 11(2):839-49, 2012.
31. Fiedler J, Etzel N and Brenner RE: To go or not to go: Migration of human mesenchymal progenitor cells stimulated by isoforms of PDGF. *J Cell Biochem* 93(5):990-8, 2004.
32. Meyer RD, Sacks DB and Rahimi N: IQGAP1-dependent signaling pathway regulates endothelial cell proliferation and angiogenesis. *PLoS One* 3(12):e3848, 2008.

33. Tomasoni S, Longaretti L, Rota C, Morigi M, Conti S, Gotti E, Capelli C, Introna M, Remuzzi G and Benigni A: Transfer of growth factor receptor mRNA via exosomes unravels the regenerative effect of mesenchymal stem cells. *Stem Cells Dev* 22(5):772-80, 2013.
34. Collino F, Deregibus MC, Bruno S, Sterpone L, Aghemo G, Viltono L, Tetta C and Camussi G: Microvesicles derived from adult human bone marrow and tissue specific mesenchymal stem cells shuttle selected pattern of miRNAs. *PLoS One* 5(7):e11803, 2010.
35. Nakamura Y, Miyaki S, Ishitobi H, Matsuyama S, Nakasa T, Kamei N, Akimoto T, Higashi Y and Ochi M: Mesenchymal-stem-cell-derived exosomes accelerate skeletal muscle regeneration. *FEBS Lett* 589(11):1257-65, 2015.
36. You L, Gu W, Chen L, Pan L, Chen J and Peng Y: MiR-378 overexpression attenuates high glucose-suppressed osteogenic differentiation through targeting CASP3 and activating PI3K/Akt signaling pathway. *Int J Clin Exp Pathol* 7(10):7249-61, 2014.
37. Guan X, Gao Y, Zhou J, Wang J, Zheng F, Guo F, Chang A, Li X and Wang B: miR-223 Regulates Adipogenic and Osteogenic Differentiation of Mesenchymal Stem Cells Through a C/EBPs/miR-223/FGFR2 Regulatory Feedback Loop. *Stem Cells* 33(5):1589-600, 2015.
38. Zeng Y, Qu X, Li H, Huang S, Wang S, Xu Q, Lin R, Han Q, Li J and Zhao RC: MicroRNA-100 regulates osteogenic differentiation of human adipose-derived mesenchymal stem cells by targeting BMP2. *FEBS Lett* 586(16):2375-81, 2012.
39. Krist B, Florczyk U, Pietraszek-Gremplewicz K, Jozkowicz A and Dulak J: The Role of miR-378a in Metabolism, Angiogenesis, and Muscle Biology. *Int J Endocrinol* 2015:281756, 2015.
40. Shi L, Fisslthaler B, Zippel N, Fromel T, Hu J, Elgheznawy A, Heide H, Popp R and Fleming I: MicroRNA-223 antagonizes angiogenesis by targeting beta1 integrin and preventing growth factor signaling in endothelial cells. *Circ Res* 113(12):1320-30, 2013.
41. Dignat-George F and Boulanger CM: The many faces of endothelial microparticles. *Arterioscler Thromb Vasc Biol* 31(1):27-33, 2011.
42. Curtis AM, Wilkinson PF, Gui M, Gales TL, Hu E and Edelberg JM: p38 mitogen-activated protein kinase targets the production of proinflammatory endothelial microparticles. *J Thromb Haemost* 7(4):701-9, 2009.
43. Leroyer AS, Anfosso F, Lacroix R, Sabatier F, Simoncini S, Njock SM, Jourde N, Brunet P, Camoin-Jau L, Sampol J, et al.: Endothelial-derived microparticles: Biological conveyors at the crossroad of inflammation, thrombosis and angiogenesis. *Thromb Haemost* 104(3):456-63, 2010.
44. Nomura S and Shimizu M: Clinical significance of procoagulant microparticles. *J Intensive Care* 3(1):2, 2015.
45. Jy W, Jimenez JJ, Mauro LM, Horstman LL, Cheng P, Ahn ER, Bidot CJ and Ahn YS: Endothelial microparticles induce formation of platelet aggregates via a von Willebrand factor/ristocetin dependent pathway, rendering them resistant to dissociation. *J Thromb Haemost* 3(6):1301-8, 2005.
46. Zhang Y, Meng H, Ma R, He Z, Wu X, Cao M, Yao Z, Zhao L, Li T, Deng R, et al.: Circulating Microparticles, Blood Cells, and Endothelium Induce Procoagulant Activity in Sepsis through Phosphatidylserine Exposure. *Shock* 45(3):299-307, 2016.
47. Andrews AM and Rizzo V: Microparticle-Induced Activation of the Vascular Endothelium Requires Caveolin-1/Caveolae. *PLoS One* 11(2):e0149272, 2016.

48. Deregibus MC, Cantaluppi V, Calogero R, Lo Iacono M, Tetta C, Biancone L, Bruno S, Bussolati B and Camussi G: Endothelial progenitor cell derived microvesicles activate an angiogenic program in endothelial cells by a horizontal transfer of mRNA. *Blood* 110(7):2440-8, 2007.
49. Jansen F, Yang X, Hoelscher M, Cattelan A, Schmitz T, Proebsting S, Wenzel D, Vosen S, Franklin BS, Fleischmann BK, et al.: Endothelial microparticle-mediated transfer of MicroRNA-126 promotes vascular endothelial cell repair via SPRED1 and is abrogated in glucose-damaged endothelial microparticles. *Circulation* 128(18):2026-38, 2013.
50. Taraboletti G, D'Ascenzo S, Borsotti P, Giavazzi R, Pavan A and Dolo V: Shedding of the matrix metalloproteinases MMP-2, MMP-9, and MT1-MMP as membrane vesicle-associated components by endothelial cells. *Am J Pathol* 160(2):673-80, 2002.
51. Lacroix R, Sabatier F, Mialhe A, Basire A, Pannell R, Borghi H, Robert S, Lamy E, Plawinski L, Camoin-Jau L, et al.: Activation of plasminogen into plasmin at the surface of endothelial microparticles: a mechanism that modulates angiogenic properties of endothelial progenitor cells in vitro. *Blood* 110(7):2432-9, 2007.
52. Wu Y and Madri J: Insights into monocyte-driven osteoclastogenesis and its link with hematopoiesis: regulatory roles of PECAM-1 (CD31) and SHP-1. *Crit Rev Immunol* 30(5):423-33, 2010.
53. Alique M, Ruiz-Torres MP, Bodega G, Noci MV, Troyano N, Bohorquez L, Luna C, Luque R, Carmona A, Carracedo J, et al.: Microvesicles from the plasma of elderly subjects and from senescent endothelial cells promote vascular calcification. *Aging (Albany NY)* 9(3):778-789, 2017.
54. Buendia P, Montes de Oca A, Madueno JA, Merino A, Martin-Malo A, Aljama P, Ramirez R, Rodriguez M and Carracedo J: Endothelial microparticles mediate inflammation-induced vascular calcification. *FASEB J* 29(1):173-81, 2015.
55. Kagiya T and Nakamura S: Expression profiling of microRNAs in RAW264.7 cells treated with a combination of tumor necrosis factor alpha and RANKL during osteoclast differentiation. *J Periodontal Res* 48(3):373-85, 2013.
56. Huynh N, VonMoss L, Smith D, Rahman I, Felemban MF, Zuo J, Rody WJ, Jr., McHugh KP and Holliday LS: Characterization of Regulatory Extracellular Vesicles from Osteoclasts. *J Dent Res*, 2016.
57. Ismail N, Wang Y, Dakhllallah D, Moldovan L, Agarwal K, Batte K, Shah P, Wisler J, Eubank TD, Tridandapani S, et al.: Macrophage microvesicles induce macrophage differentiation and miR-223 transfer. *Blood* 121(6):984-95, 2013.
58. Ekstrom K, Omar O, Graneli C, Wang X, Vazirisani F and Thomsen P: Monocyte exosomes stimulate the osteogenic gene expression of mesenchymal stem cells. *PLoS One* 8(9):e75227, 2013.
59. Essayagh S, Xuereb JM, Terrisse AD, Tellier-Cirioni L, Pipy B and Sie P: Microparticles from apoptotic monocytes induce transient platelet recruitment and tissue factor expression by cultured human vascular endothelial cells via a redox-sensitive mechanism. *Thromb Haemost* 98(4):831-7, 2007.
60. Del Conde I, Shrimpton CN, Thiagarajan P and Lopez JA: Tissue-factor-bearing microvesicles arise from lipid rafts and fuse with activated platelets to initiate coagulation. *Blood* 106(5):1604-11, 2005.

61. Li J, Zhang Y, Liu Y, Dai X, Li W, Cai X, Yin Y, Wang Q, Xue Y, Wang C, et al.: Microvesicle-mediated transfer of microRNA-150 from monocytes to endothelial cells promotes angiogenesis. *J Biol Chem* 288(32):23586-96, 2013.
62. Johnson BL, Iii, Kuethe JW and Caldwell CC: Neutrophil derived microvesicles: emerging role of a key mediator to the immune response. *Endocr Metab Immune Disord Drug Targets* 14(3):210-7, 2014.
63. Prakash PS, Caldwell CC, Lentsch AB, Pritts TA and Robinson BR: Human microparticles generated during sepsis in patients with critical illness are neutrophil-derived and modulate the immune response. *J Trauma Acute Care Surg* 73(2):401-6; discussion 406-7, 2012.
64. Timar CI, Lorincz AM, Csepanyi-Komi R, Valyi-Nagy A, Nagy G, Buzas EI, Ivanyi Z, Kittel A, Powell DW, McLeish KR, et al.: Antibacterial effect of microvesicles released from human neutrophilic granulocytes. *Blood* 121(3):510-8, 2013.
65. Pluskota E, Woody NM, Szpak D, Ballantyne CM, Soloviev DA, Simon DI and Plow EF: Expression, activation, and function of integrin alphaMbeta2 (Mac-1) on neutrophil-derived microparticles. *Blood* 112(6):2327-35, 2008.
66. Koppler B, Cohen C, Schlondorff D and Mack M: Differential mechanisms of microparticle transfer to B cells and monocytes: anti-inflammatory properties of microparticles. *Eur J Immunol* 36(3):648-60, 2006.
67. Nolan S, Dixon R, Norman K, Hellewell P and Ridger V: Nitric oxide regulates neutrophil migration through microparticle formation. *Am J Pathol* 172(1):265-73, 2008.
68. Pitanga TN, de Aragao Franca L, Rocha VC, Meirelles T, Borges VM, Goncalves MS, Pontes-de-Carvalho LC, Noronha-Dutra AA and dos-Santos WL: Neutrophil-derived microparticles induce myeloperoxidase-mediated damage of vascular endothelial cells. *BMC Cell Biol* 15:21, 2014.
69. Midura EF, Kuethe JW, Rice TC, Veile R, England LG, Friend LA, Caldwell CC and Goodman MD: Impact of Platelets and Platelet-Derived Microparticles on Hypercoagulability Following Burn Injury. *Shock* 45(1):82-7, 2016.
70. Iwamoto S, Kawasaki T, Kambayashi J, Ariyoshi H and Monden M: Platelet microparticles: a carrier of platelet-activating factor? *Biochem Biophys Res Commun* 218(3):940-4, 1996.
71. Sinauridze EI, Kireev DA, Popenko NY, Pichugin AV, Panteleev MA, Krymskaya OV and Ataullakhanov FI: Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thromb Haemost* 97(3):425-34, 2007.
72. Majka M, Kijowski J, Lesko E, Gozdick J, Zupanska B and Ratajczak MZ: Evidence that platelet-derived microvesicles may transfer platelet-specific immunoreactive antigens to the surface of endothelial cells and CD34+ hematopoietic stem/ progenitor cells--implication for the pathogenesis of immune thrombocytopenias. *Folia Histochem Cytobiol* 45(1):27-32, 2007.
73. Curry N, Raja A, Beavis J, Stanworth S and Harrison P: Levels of procoagulant microvesicles are elevated after traumatic injury and platelet microvesicles are negatively correlated with mortality. *J Extracell Vesicles* 3:25625, 2014.
74. Barry OP and FitzGerald GA: Mechanisms of cellular activation by platelet microparticles. *Thromb Haemost* 82(2):794-800, 1999.

75. Merten M, Pakala R, Thiagarajan P and Benedict CR: Platelet microparticles promote platelet interaction with subendothelial matrix in a glycoprotein IIb/IIIa-dependent mechanism. *Circulation* 99(19):2577-82, 1999.
76. Balvers K, Curry N, Kleinveld DJ, Boing AN, Nieuwland R, Goslings JC and Juffermans NP: Endogenous microparticles drive the proinflammatory host immune response in severely injured trauma patients. *Shock* 43(4):317-21, 2015.
77. Barry OP, Kazanietz MG, Pratico D and FitzGerald GA: Arachidonic acid in platelet microparticles up-regulates cyclooxygenase-2-dependent prostaglandin formation via a protein kinase C/mitogen-activated protein kinase-dependent pathway. *J Biol Chem* 274(11):7545-56, 1999.
78. Laffont B, Corduan A, Rousseau M, Duchez AC, Lee CH, Boilard E and Provost P: Platelet microparticles reprogram macrophage gene expression and function. *Thromb Haemost* 115(2):311-23, 2016.
79. Bei JJ, Liu C, Peng S, Liu CH, Zhao WB, Qu XL, Chen Q, Zhou Z, Yu ZP, Peter K, et al.: Staphylococcal SSL5-induced platelet microparticles provoke proinflammatory responses via the CD40/TRAF6/NFkappaB signalling pathway in monocytes. *Thromb Haemost* 115(3):632-45, 2016.
80. Jy W, Mao WW, Horstman L, Tao J and Ahn YS: Platelet microparticles bind, activate and aggregate neutrophils in vitro. *Blood Cells Mol Dis* 21(3):217-31; discussion 231a, 1995.
81. Sprague DL, Elzey BD, Crist SA, Waldschmidt TJ, Jensen RJ and Ratliff TL: Platelet-mediated modulation of adaptive immunity: unique delivery of CD154 signal by platelet-derived membrane vesicles. *Blood* 111(10):5028-36, 2008.
82. Kim HK, Song KS, Chung JH, Lee KR and Lee SN: Platelet microparticles induce angiogenesis in vitro. *Br J Haematol* 124(3):376-84, 2004.
83. Laffont B, Corduan A, Ple H, Duchez AC, Cloutier N, Boilard E and Provost P: Activated platelets can deliver mRNA regulatory Ago2*microRNA complexes to endothelial cells via microparticles. *Blood* 122(2):253-61, 2013.
84. Gruber R, Varga F, Fischer MB and Watzek G: Platelets stimulate proliferation of bone cells: involvement of platelet-derived growth factor, microparticles and membranes. *Clin Oral Implants Res* 13(5):529-35, 2002.
85. Baj-Krzyworzeka M, Majka M, Pratico D, Ratajczak J, Vilaire G, Kijowski J, Reca R, Janowska-Wieczorek A and Ratajczak MZ: Platelet-derived microparticles stimulate proliferation, survival, adhesion, and chemotaxis of hematopoietic cells. *Exp Hematol* 30(5):450-9, 2002.
86. Chung SM, Bae ON, Lim KM, Noh JY, Lee MY, Jung YS and Chung JH: Lysophosphatidic acid induces thrombogenic activity through phosphatidylserine exposure and procoagulant microvesicle generation in human erythrocytes. *Arterioscler Thromb Vasc Biol* 27(2):414-21, 2007.
87. Llorente-Cortes V, Otero-Vinas M, Camino-Lopez S, Llampayas O and Badimon L: Aggregated low-density lipoprotein uptake induces membrane tissue factor procoagulant activity and microparticle release in human vascular smooth muscle cells. *Circulation* 110(4):452-9, 2004.
88. Nahar NN, Missana LR, Garimella R, Tague SE and Anderson HC: Matrix vesicles are carriers of bone morphogenetic proteins (BMPs), vascular endothelial growth factor (VEGF), and noncollagenous matrix proteins. *J Bone Miner Metab* 26(5):514-9, 2008.

89. Cooper LF, Yliheikkilä PK, Felton DA and Whitson SW: Spatiotemporal assessment of fetal bovine osteoblast culture differentiation indicates a role for BSP in promoting differentiation. *J Bone Miner Res* 13(4):620-32, 1998.
90. Gordon JA, Tye CE, Sampaio AV, Underhill TM, Hunter GK and Goldberg HA: Bone sialoprotein expression enhances osteoblast differentiation and matrix mineralization in vitro. *Bone* 41(3):462-73, 2007.
91. Wang J, Zhou HY, Salih E, Xu L, Wunderlich L, Gu X, Hofstaetter JG, Torres M and Glimcher MJ: Site-specific in vivo calcification and osteogenesis stimulated by bone sialoprotein. *Calcif Tissue Int* 79(3):179-89, 2006.
92. Gerber HP, Vu TH, Ryan AM, Kowalski J, Werb Z and Ferrara N: VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. *Nat Med* 5(6):623-8, 1999.
93. Horstman LL, Jy W, Jimenez JJ, Bidot C and Ahn YS: New horizons in the analysis of circulating cell-derived microparticles. *Keio J Med* 53(4):210-30, 2004.
94. Zwaal RF and Schroit AJ: Pathophysiologic implications of membrane phospholipid asymmetry in blood cells. *Blood* 89(4):1121-32, 1997.
95. Lacroix R, Plawinski L, Robert S, Dœuvre L, Sabatier F, Martinez de Lizarrondo S, Mezzapesa A, Anfosso F, Leroyer AS, Poullin P, et al.: Leukocyte- and endothelial-derived microparticles: a circulating source for fibrinolysis. *Haematologica* 97(12):1864-72, 2012.
96. Hong Y, Eleftheriou D, Hussain AA, Price-Kuehne FE, Savage CO, Jayne D, Little MA, Salama AD, Klein NJ and Brogan PA: Anti-neutrophil cytoplasmic antibodies stimulate release of neutrophil microparticles. *J Am Soc Nephrol* 23(1):49-62, 2012.
97. Dalli J, Norling LV, Renshaw D, Cooper D, Leung KY and Perretti M: Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. *Blood* 112(6):2512-9, 2008.
98. Eken C, Martin PJ, Sadallah S, Treves S, Schaller M and Schifferli JA: Ectosomes released by polymorphonuclear neutrophils induce a MerTK-dependent anti-inflammatory pathway in macrophages. *J Biol Chem* 285(51):39914-21, 2010.
99. Woodfin A, Voisin MB and Nourshargh S: PECAM-1: a multi-functional molecule in inflammation and vascular biology. *Arterioscler Thromb Vasc Biol* 27(12):2514-23, 2007.
100. Cantaluppi V, Gatti S, Medica D, Figliolini F, Bruno S, Deregibus MC, Sordi A, Biancone L, Tetta C and Camussi G: Microvesicles derived from endothelial progenitor cells protect the kidney from ischemia-reperfusion injury by microRNA-dependent reprogramming of resident renal cells. *Kidney Int* 82(4):412-27, 2012.
101. Lee DY, Deng Z, Wang CH and Yang BB: MicroRNA-378 promotes cell survival, tumor growth, and angiogenesis by targeting SuFu and Fus-1 expression. *Proc Natl Acad Sci U S A* 104(51):20350-5, 2007.
102. Zhang J, Guan J, Niu X, Hu G, Guo S, Li Q, Xie Z, Zhang C and Wang Y: Exosomes released from human induced pluripotent stem cells-derived MSCs facilitate cutaneous wound healing by promoting collagen synthesis and angiogenesis. *J Transl Med* 13:49, 2015.
103. Shabbir A, Cox A, Rodriguez-Menocal L, Salgado M and Van Badiavas E: Mesenchymal Stem Cell Exosomes Induce Proliferation and Migration of Normal and Chronic Wound Fibroblasts, and Enhance Angiogenesis In Vitro. *Stem Cells Dev* 24(14):1635-47, 2015.

104. Wu Y, Tworkoski K, Michaud M and Madri JA: Bone marrow monocyte PECAM-1 deficiency elicits increased osteoclastogenesis resulting in trabecular bone loss. *J Immunol* 182(5):2672-9, 2009.
105. Shapiro IM, Landis WJ and Risbud MV: Matrix vesicles: Are they anchored exosomes? *Bone* 79:29-36, 2015.
106. Keerthikumar S, Chisanga D, Ariyaratne D, Al Saffar H, Anand S, Zhao K, Samuel M, Pathan M, Jois M, Chilamkurti N, et al.: ExoCarta: A Web-Based Compendium of Exosomal Cargo. *J Mol Biol* 428(4):688-92, 2016.
107. Kim SJ, Moon GJ and Bang OY: Biomarkers for stroke. *J Stroke* 15(1):27-37, 2013.
108. Berezin A, Zulli A, Kerrigan S, Petrovic D and Kruzliak P: Predictive role of circulating endothelial-derived microparticles in cardiovascular diseases. *Clin Biochem* 48(9):562-8, 2015.
109. Dumache R, Rogobete AF, Bedreag OH, Sarandan M, Cradigati AC, Papurica M, Dumbuleu CM, Nartita R and Sandesc D: Use of miRNAs as biomarkers in sepsis. *Anal Cell Pathol (Amst)* 2015:186716, 2015.
110. Vinuela-Berni V, Doniz-Padilla L, Figueroa-Vega N, Portillo-Salazar H, Abud-Mendoza C, Baranda L and Gonzalez-Amaro R: Proportions of several types of plasma and urine microparticles are increased in patients with rheumatoid arthritis with active disease. *Clin Exp Immunol* 180(3):442-51, 2015.
111. Murata K, Ito H, Yoshitomi H, Yamamoto K, Fukuda A, Yoshikawa J, Furu M, Ishikawa M, Shibuya H and Matsuda S: Inhibition of miR-92a enhances fracture healing via promoting angiogenesis in a model of stabilized fracture in young mice. *J Bone Miner Res* 29(2):316-26, 2014.
112. Dai S, Wei D, Wu Z, Zhou X, Wei X, Huang H and Li G: Phase I clinical trial of autologous ascites-derived exosomes combined with GM-CSF for colorectal cancer. *Mol Ther* 16(4):782-90, 2008.
113. Zhu YG, Feng XM, Abbott J, Fang XH, Hao Q, Monsel A, Qu JM, Matthay MA and Lee JW: Human mesenchymal stem cell microvesicles for treatment of Escherichia coli endotoxin-induced acute lung injury in mice. *Stem Cells* 32(1):116-25, 2014.
114. Lai RC, Arslan F, Lee MM, Sze NS, Choo A, Chen TS, Salto-Tellez M, Timmers L, Lee CN, El Oakley RM, et al.: Exosome secreted by MSC reduces myocardial ischemia/reperfusion injury. *Stem Cell Res* 4(3):214-22, 2010.
115. Dorronsoro A and Robbins PD: Regenerating the injured kidney with human umbilical cord mesenchymal stem cell-derived exosomes. *Stem Cell Res Ther* 4(2):39, 2013.
116. Moest T, Koehler F, Prechtel C, Schmitt C, Watzek G and Schlegel KA: Bone formation in peri-implant defects grafted with microparticles: a pilot animal experimental study. *J Clin Periodontol* 41(10):990-8, 2014.
117. O'Brien K, Lowry MC, Corcoran C, Martinez VG, Daly M, Rani S, Gallagher WM, Radomski MW, MacLeod RA and O'Driscoll L: miR-134 in extracellular vesicles reduces triple-negative breast cancer aggression and increases drug sensitivity. *Oncotarget* 6(32):32774-89, 2015.
118. Fais S, O'Driscoll L, Borrás FE, Buzas E, Camussi G, Cappello F, Carvalho J, Cordeiro da Silva A, Del Portillo H, El Andaloussi S, et al.: Evidence-Based Clinical Use of Nanoscale Extracellular Vesicles in Nanomedicine. *ACS Nano*, 2016.
119. Link DP, van den Dolder J, van den Beucken JJ, Wolke JG, Mikos AG and Jansen JA: Bone response and mechanical strength of rabbit femoral defects filled with injectable CaP cements containing TGF-beta 1 loaded gelatin microparticles. *Biomaterials* 29(6):675-82, 2008.

120. Dang PN, Dwivedi N, Phillips LM, Yu X, Herberg S, Bowerman C, Solorio LD, Murphy WL and Alsberg E: Controlled Dual Growth Factor Delivery From Microparticles Incorporated Within Human Bone Marrow-Derived Mesenchymal Stem Cell Aggregates for Enhanced Bone Tissue Engineering via Endochondral Ossification. *Stem Cells Transl Med* 5(2):206-17, 2016.
121. Piao ZG, Kim JS, Son JS, Lee SY, Fang XH, Oh JS, You JS and Kim SG: Osteogenic evaluation of collagen membrane containing drug-loaded polymeric microparticles in a rat calvarial defect model. *Tissue Eng Part A* 20(23-24):3322-31, 2014.
122. Jing D, Hao J, Shen Y, Tang G, Li ML, Huang SH and Zhao ZH: The role of microRNAs in bone remodeling. *Int J Oral Sci* 7(3):131-43, 2015.
123. Svensson D, Gidlof O, Turczynska KM, Erlinge D, Albinsson S and Nilsson BO: Inhibition of microRNA-125a promotes human endothelial cell proliferation and viability through an antiapoptotic mechanism. *J Vasc Res* 51(3):239-45, 2014.
124. Guo LJ, Liao L, Yang L, Li Y and Jiang TJ: MiR-125a TNF receptor-associated factor 6 to inhibit osteoclastogenesis. *Exp Cell Res* 321(2):142-52, 2014.
125. Wen P, Cao H, Fang L, Ye H, Zhou Y, Jiang L, Su W, Xu H, He W, Dai C, et al.: miR-125b/Ets1 axis regulates transdifferentiation and calcification of vascular smooth muscle cells in a high-phosphate environment. *Exp Cell Res* 322(2):302-12, 2014.
126. Schmidt Y, Simunovic F, Strassburg S, Pfeifer D, Stark GB and Finkenzeller G: miR-126 regulates platelet-derived growth factor receptor- α expression and migration of primary human osteoblasts. *Biol Chem* 396(1):61-70, 2015.
127. Huang F, Fang ZF, Hu XQ, Tang L, Zhou SH and Huang JP: Overexpression of miR-126 promotes the differentiation of mesenchymal stem cells toward endothelial cells via activation of PI3K/Akt and MAPK/ERK pathways and release of paracrine factors. *Biol Chem* 394(9):1223-33, 2013.
128. Zhang Y, Liu D, Chen X, Li J, Li L, Bian Z, Sun F, Lu J, Yin Y, Cai X, et al.: Secreted monocytic miR-150 enhances targeted endothelial cell migration. *Mol Cell* 39(1):133-44, 2010.
129. Huang J, Zhao L, Xing L and Chen D: MicroRNA-204 regulates Runx2 protein expression and mesenchymal progenitor cell differentiation. *Stem Cells* 28(2):357-64, 2010.
130. Tang PF, Xiong Q, Ge W and Zhang LH: The role of MicroRNAs in Osteoclasts and Osteoporosis. *Rna Biology* 11(11):1355-1363, 2014.
131. Kagiya T and Taira M: Expression of MicroRNAs in the Extracellular Microvesicles of Murine Osteoclasts. *Journal of Oral Tissue Engineering* 10(3):142-150, 2013.
132. Lou YL, Guo F, Liu F, Gao FL, Zhang PQ, Niu X, Guo SC, Yin JH, Wang Y and Deng ZF: miR-210 activates notch signaling pathway in angiogenesis induced by cerebral ischemia. *Mol Cell Biochem* 370(1-2):45-51, 2012.
133. Li Z, Meng D, Li G, Xu J, Tian K and Li Y: Overexpression of microRNA210 promotes chondrocyte proliferation and extracellular matrix deposition by targeting HIF3 α in osteoarthritis. *Mol Med Rep*, 2016.
134. Xue Y, Wei Z, Ding H, Wang Q, Zhou Z, Zheng S, Zhang Y, Hou D, Liu Y, Zen K, et al.: MicroRNA-19b/221/222 induces endothelial cell dysfunction via suppression of PGC-1 α in the progression of atherosclerosis. *Atherosclerosis* 241(2):671-81, 2015.

135. Zhan C, Ma CB, Yuan HM, Cao BY and Zhu JJ: Macrophage-derived microvesicles promote proliferation and migration of Schwann cell on peripheral nerve repair. *Biochem Biophys Res Commun* 468(1-2):343-8, 2015.

ACCEPTED

Figure 1. MV-originated miRNAs related to osteoblastic functions. Runt-related transcription factor 2 (RUNX2), transforming growth factor-β(TGF-β), bone morphogenetic protein (BMP), phosphatidylinositol-3 kinase (PI3K), nuclear factor kappa B (NF-κB), histone deacetylases (HDAC), mitogen-activated protein kinase (MAPK), mechanistic target of rapamycin (mTOR)



Figure 2. MV-originated miRNAs related to endothelial cell functions.

Runt-related transcription factor 2 (RUNX2), transforming growth factor(TGF), bone morphogenetic protein (BMP), phosphatidylinositol-3 kinase (PI3K), nuclear factor kappa B (NF-κB), histone deacetylases (HDAC), mitogen-activated protein kinase (MAPK)

extracellular signal–regulated kinases (ERK), mechanistic target of rapamycin (mTOR), Insulin-like growth factor 1 (IGF-1) platelet-derived growth factor receptor (PDGFR) Semaphorin-6A (SEMA6A) Phosphatase and tensin homolog (PTEN) Hypoxia-inducible factor 1-alpha (HIF-1α) vascular endothelial growth factor (VEGF)




Figure 3. Known MV-based crosstalk mechanisms during fracture healing.

Abbreviations: anionic phospholipid (AP), tissue factor (TF), arachidonic acid (AA), platelet-activating factor (PAF), P-selectin glycoprotein ligand-1 (PSGL-1), platelet-derived growth factor receptor (PDGFR), epidermal growth factor receptor (EGFR), mitogen-activated protein kinase (MAPK), phosphatidylinositol-3 kinase (PI-3K), Signal transducer and activator of transcription (STAT), argonaute 2 (Ago2), endothelial protein C receptor (EPCR), thrombomodulin (TM), von Willebrand factor (vWf), plasminogen activator (PA), phosphatidylserines (PS), platelet endothelial cell adhesion molecule (PECAM), intercellular adhesion molecule (ICAM), vascular cell adhesion molecule (VCAM), bone morphogenetic protein (BMP), ezrin (EZR), IQ motif containing GTPase activating protein 1 (IQGAP-1), insulin-like growth factor-1 receptor (IGF-1R), stromal cell-derived factor 1 (SDF1), growth factor (GF), receptor activator of nuclear factor kappa-B ligand (RANKL)




Table 1. summary of EV-based miRNAs related to osteogenesis

miRNA type	Recipient cell	Donor cells	References
miR-100	MSC	MSC	(122), (34), (38)
miR-125a	endothelial cell, osteoclast	MSC	(34), (123), (124)
miR-125b	osteoblast	MSC	(122), (34), (125),
miR-126	endothelial cell, MSC, osteoblast	endothelial cell	(49), (126), (127)
miR-133a	osteoblast	MSC	(34), (122)
miR-135a	osteoblast	MSC	(34), (122)
miR-137	osteoblast	MSC	(34), (122)
miR-150	endothelial cell	monocyte	(61)
miR-155	endothelial cell	osteoclast, monocyte	(55), (128)
miR-204	MSC	MSC	(34), (129)
miR-205	osteoblast	MSC	(34)
miR-21	osteoblast, chondrocyte, endothelial cell,	endothelial, osteoclast	(130), (55), (122), (131)

	MSC, osteoclast		
miR-210	osteoblast, chondrocyte, endothelial cell, MSC	osteoclast	(131), (132), (122), (133)
miR-217	osteoblast	MSC	(34), (122)
miR-221	endothelial cell	MSC	(34), (134)
miR-223	osteoblast,osteoclast, endothelial cell	MSC, osteoclast, macrophage	(34), (122), (135), (130), (55), (131)
miR-23a	osteoblast, MSC	MSC	(34), (122)
miR-24	osteoblast	MSC	(34), (122)
miR-30	osteoblast	MSC	(34), (122)
miR-31	endothelial cell, osteoclast, osteoblast	MSC	(34), (122)
miR-338	osteoblast	MSC	(34), (122)
miR-34	osteoblast	MSC	(34), (122)
miR-378	osteoclast, endothelial cell	osteoclast	(36, 55, 130),
miR-451	endothelial cell	MSC	(34)